


AFATL-TR-87-61

DTIC FILE COPY

**Synthesis of 1,4-Butanediamine-1,
1,4,4-d₄, 1,4-Butanediamine-2,
2,3,3-d₄ and Their Respective
BIS (Ammonium Nitrate) Salts**

Dr Alan P Marchand
Dr N Satyanarayana
Dr Robert L McKenney Jr
Mr Stephen R Struck

NORTH TEXAS STATE UNIV
DEPARTMENT OF CHEMISTRY
NT STATION, BOX 5068
DENTON, TEXAS 76203-5068



DECEMBER 1987

INTERIM REPORT FOR PERIOD OCTOBER 1985-OCTOBER 1987

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

AIR FORCE ARMAMENT LABORATORY

Air Force Systems Command ■ United States Air Force ■ Eglin Air Force Base, Florida

88 1 12 164

AD-A194 518

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-018	
1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) N/A			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFATL-TR-87-61		
6a. NAME OF PERFORMING ORGANIZATION North Texas State University		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Munitions Division		
6c. ADDRESS (City, State, and ZIP Code) Dept of Chemistry, North Texas State University NT Station, Box 5068 Denton TX 76203-5068			7b. ADDRESS (City, State, and ZIP Code) Air Force Armament Laboratory Eglin AFB FL 32542-5434		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Munitions Division		8b. OFFICE SYMBOL (If applicable) AFATL/MN	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F08635-86-K-0078 F08635-87-K-0101		
8c. ADDRESS (City, State, and ZIP Code) Air Force Armament Laboratory Eglin AFB FL 32542-5434			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 62602F	PROJECT NO. 2502	TASK NO. 10
					WORK UNIT ACCESSION 12, 13,
11. TITLE (Include Security Classification) Synthesis of 1,4-Butanediamine-1,1,4,4-d ₄ , 1,4-Butanediamine-2,2,3,3-d ₄ and Their Respective Bis(Ammonium Nitrate) Salts					
12. PERSONAL AUTHOR(S) A.P. Marchand, N. Satyanarayana, R.L. McKenney Jr., S.R. Struck					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM Oct 85 TO Oct 87		14. DATE OF REPORT (Year, Month, Day) December 1987	
15. PAGE COUNT 16					
16. SUPPLEMENTARY NOTATION Availability of report specified on verso of front cover					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
07	02		Synthesis, Site-specified ² H Labelled Putrescines, Site-specific ² H Labelled Putrescinium Nitrate Salts		
07	03				
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
Synthesis of the title compounds from diethyl succinate and from dimethyl acetylenedicarboxylate in 16 percent overall yield, respectively, and their bis(ammonium nitrate) salts in 62 and 60 percent yields, respectively, are described. Physical properties and nuclear magnetic resonance, infrared and mass spectral data are presented.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Robert L. McKenney Jr			22b. TELEPHONE (Include Area Code) 904 882-4212		22c. OFFICE SYMBOL AFATL/MNE

PREFACE

This report describes the work performed by personnel at the Department of Chemistry, North Texas State University, NT Station, Box 5063, Denton TX 76203-5068 under contracts F08635-86-K-0078 and F08635-87-K-0101 during the period January 1986-September 1987 and by personnel at the Energetic Materials Branch (MNE), Munitions Division (MN), Air Force Armament Laboratory (AFATL), Armament Division (AD), Eglin AFB FL 32542-5434, under project 2502, "Conventional Ordnance Technology", Task 10, "Advanced Explosives", Work Unit 12, "Explosive Response to Thermal and Shock Stimuli" during the time period of October 1985 to October 1987. The contract program manager was Dr Alan P. Marchand and the AFATL in-house program manager was Dr Robert L. McKenney Jr.

This report is one of several planned reports that will describe the synthesis and complete thermal characterization of specific organodi- and polyammonium nitrate salts. Other reports will follow at appropriate times during the progress of this effort.



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A-1	

TABLE OF CONTENTS

Section	Title	Page
I	INTRODUCTION.	1
II	DISCUSSION.	2
III	EXPERIMENTAL.	4
IV	CONCLUSIONS	7
	REFERENCES.	9

LIST OF FIGURES

Figure	Title	Page
1	Summary of Reaction Schemes (I and II) for the Preparation of 1,4-Butanediamine-1,1,4,4-d ₄ (1a) and 1,4-Butanediamine-2,2,3,3-d ₄ (1b)	3

SECTION I

INTRODUCTION

There is considerable current interest in the study of kinetic isotope effects in thermal decompositions of explosives.¹⁻³ As part of a study of deuterium isotope effects on the decomposition of 1,4-butanediammonium dinitrate (BDD), it was necessary to synthesize 1,4-butanediamine-1,1,4,4-d₄ (putrescine-1,1,4,4-d₄ **1a**), the corresponding 2,2,3,3-d₄ isomer (putrescine-2,2,3,3-d₄, **1b**) and the bis(ammonium nitrate) salts of **1a** and **1b** (i.e., **1c** and **1d**, respectively). The synthesis of the bis(ammonium acetate) salt of **1a** via lithium aluminum deuteride reduction of succinonitrile has been reported by Mayerl and Hesse;⁴ however, their reported yield was only ca. 1.4%. Our own experience with this reaction similarly produced **1a** in disappointingly low yield. In contrast to this result, deuterogenation of succinonitrile over platinum oxide catalyst in aqueous ethanolic (C₂H₅OD) hydrochloric acid (DCl) solution has been reported to afford 1,4-butanediamine-1,1,4,4-d₄ dihydrochloride in 40% yield.⁵

It is most advantageous to prepare BDD directly via reaction of putrescine with nitric acid. It is not convenient to liberate the free amine from one of its free bases directly.

SECTION II


DISCUSSION

To synthesize **1a**, we employed a Gabriel synthesis⁶ as outlined in Scheme I of Figure 1. Lithium aluminum deuteride reduction of diethyl succinate (**2**) afforded 1,4-butanediol-1,1,4,4-d₄ (**3**) in 66% yield. Reaction of **3** with phosphorus tribromide afforded the corresponding dibromide (**4**, 60%). Subsequent reaction of **4** with two equivalents of potassium phthalimide afforded the corresponding diphthalimido derivative (**5**, 88%). Finally, pure **1a** was produced in 47% yield (16% overall from diethyl succinate) by refluxing a solution of **5** in absolute ethanol with excess hydrazine hydrate.

Our corresponding synthesis of **1b** is summarized in Scheme II of Figure 1. Dimethyl acetylenedicarboxylate (**6**) was deuterogenated with deuterium gas over 5% palladized charcoal catalyst in methanol, thereby affording dimethyl succinate-2,2,3,3-d₄ (**7**, 35%). Lithium aluminum hydride reduction of **7** afforded 1,4-butanediol-2,2,3,3-d₄ (**8**, 85%). Thereafter, conversion of **8** to **1b** followed the steps outlined in Scheme I (Figure 1) for the conversion of **3** to **1a**. Application of this reaction sequence afforded **1b** in 18% overall yield.

The conversion of **1a** and **1b** to their respective bis(ammonium nitrate) salts (**1c** and **1d**, respectively) was effected in 62% and 60% yield, respectively, via reaction of each diamine with concentrated nitric acid.⁷ The Mass spectra of **1c** and **1d** were obtained by using the direct inlet solids probe technique at a temperature slightly above their melting points. Analysis of the fragmentation patterns thereby obtained suggests that in each case the salt undergoes initial decomposition into the free diamine and nitric acid.

Scheme I



Scheme II




Figure 1. Summary of Reaction Schemes (I and II) for the Preparation of 1,4-Butanediamine-1,1,4,4-d₄ (1a) and 1,4-Butanediamine-2,2,3,3-d₄ (1b).

SECTION III

EXPERIMENTAL

Melting points and boiling points are uncorrected. Proton NMR spectra of **1a** and **1b** were obtained on a Hitachi-Perkin Elmer Model R-243 NMR spectrometer (60 MHz) and by using a JEOL FX-90Q FT-NMR spectrometer (90 MHz). Proton NMR spectra of **1c** and **1d** were obtained on a Varian EM-390 spectrometer (90 MHz). Signals are reported in ppm (δ) downfield from internal tetramethylsilane for **1a** and **1b** and downfield from internal sodium 3-(trimethylsilyl)propionate-2,2,3,3- d_4 for **1c** and **1d**. Infrared spectra of **1a** and **1b** were obtained on a Perkin-Elmer Model 1330 IR spectrophotometer. Infrared spectra of **1c** and **1d** were obtained on a Mattson Cygnus 25 FTIR spectrometer. Mass spectra of **1a** and **1b** were obtained by using a Hewlett-Packard Model 5970A GC/MS system operating at 70eV. Mass spectra of **1c** and **1d** were obtained on a Finnigan MAT Model 5100 GC/MS system operating at 70 eV.

1,4-Butanediol-1,1,4,4- d_4 (3). A solution of **2** (13.92 g, 80 mmol) in dry tetrahydrofuran (THF, 100 ml) was added dropwise to a stirred slurry of lithium aluminum deuteride (10.08 g, 240 mmol) in dry THF (100 ml). The reaction mixture was refluxed for 4 h, at which time the reaction mixture was cooled and then quenched via successive addition of water (10 ml), 10% aqueous sodium hydroxide solution (10 ml), and water (30 ml). The resulting mixture was then filtered to remove precipitated aluminum salts. The filtrate was dried (anhydrous magnesium sulfate) and filtered, and the filtrate was concentrated in vacuo to afford crude **3** (5.1 g). The crude product was purified via column chromatography (silica gel absorbent, ethyl acetate eluent). The material thereby obtained was distilled in vacuo to afford pure **3** (5.0 g, 66%): bp 80 °C (0.5 mm). Upon cooling, the distillate solidified to afford colorless needles: mp 15 °C, (lit⁸ bp 127 °C (20 mm), mp 16 °C). IR (neat) 3300 (vs), 2095 (s), 2005 (s), 950 cm^{-1} (s); ^1H NMR (CDCl_3) δ 1.65 (br s, 4H), 3.68 (br s, 2H); mass spectrum (70 eV), m/e (relative intensity) (no molecular ion), 76 (1.7), 75 (15.4), 58 (14.1), 46 (15.8), 45 (73.0), 44 (100.0), 43 (35.7), 42 (13.6), 33 (98.3), 32 (23.7), 31 (13.6).

1,4-dibromobutane-1,1,4,4- d_4 (4). Phosphorous tribromide (14.14 g, 52 mmol) was added dropwise to cooled **3** (4.9 g, 52 mmol); the reaction mixture was cooled via application of an external ice bath. After the addition of PBr_3 had been completed, the ice bath was removed, and the reaction mixture

was stirred at ambient temperature for 13 h. The reaction mixture was then poured into water and extracted with chloroform. The organic layer was washed successively with water, aqueous sodium bicarbonate solution, and brine. The organic layer was then dried (anhydrous magnesium sulfate) and filtered, and the filtrate was concentrated in vacuo. The residue was distilled in vacuo to afford **4** as a colorless oil (7.0 g, 60%): bp 115 °C (290 mm), [lit.⁹ bp 197 °C (1 atm)]; IR (neat) 2155 (s), 660 cm⁻¹ (s); ¹H NMR (CDCl₃) δ 2.04 (s); mass spectrum (70 eV), m/e (relative intensity) 220 (molecular ion, 0.6), 141 (64.7), 139 (67.5), 109 (14.1), 81 (18.4), 79 (16.7), 59 (100.0), 43 (25.6), 41 (27.1), 40 (27.1), 32 (26.2), 31 (29.6).

1,4-Diphthalimidobutane-1,1,4,4-d₄ (5). A mixture of **4** (4.35 g, 20 mmol), potassium phthalimide (13.35 g, 72 mmol) and dimethylformamide (DMF, 60 ml) was heated on a steam bath for 4 h. The reaction mixture was then concentrated in vacuo, and the residue was poured over crushed ice. The crude product was collected via suction filtration; the residue was washed with water and then air-dried. The crude product was extracted with boiling carbon disulfide to remove N-(4-bromobutyl)phthalimide. Pure **5** (6.9 g, 38%) was thereby obtained as a colorless microcrystalline solid: mp 226 °C, (lit.⁵ mp 230 °C); IR (KBr) 3065 (m), 3025 (m), 2175 (m), 2125 (m), 1700 (br, vs), 1600 (s), 900 (s), 800 cm⁻¹ (s); ¹H NMR (CDCl₃) δ 1.70 (s, 4H), 7.61-7.86 (m, 8H); mass spectrum (70 eV), m/e (relative intensity) 352 (molecular ion, 6.8), 205 (14.1), 204 (18.3), 190 (25.7), 163 (13.3), 162 (100.0), 134 (15.7), 130 (15.7), 106 (14.7), 104 (14.1), 78 (18.3), 76 (17.8).

1,4-Butanediamine-1,1,4,4-d₄ (1a). A mixture of **5** (6.5 g, 18.5 mmol), hydrazine hydrate (2.22 g, 44.4 mmol), and absolute ethanol (120 ml) was refluxed for 4 h under nitrogen. The reaction mixture was cooled and then filtered to remove phthalhydrazide. The filtrate was dried (anhydrous magnesium sulfate) and filtered, and the filtrate was concentrated in vacuo. The residue was distilled in vacuo to afford **1a** (0.8 g, 47%) as a colorless oil: bp 85 °C (290 mm); IR (neat) 3300 (br, s), 2195 (s), 2095 (s), 1570 cm⁻¹ (br, s); ¹H NMR (CDCl₃) δ 1.38 (s, 4H), 1.52 (s, 4H); mass spectrum (70 eV), m/e (relative intensity) 92 (molecular ion, 0.1), 75 (0.4), 58 (0.6), 44 (17.0), 32 (100).

Dimethyl Succinate-2,2,3,3-d₄ (7). A stirred mixture of acetylenedicarboxylate (**6**, 10.0 g, 70.4 mmol) and 5% palladized charcoal (1.0 g) in methanol (400 ml) was purged with nitrogen gas. A balloon containing

excess deuterium gas was connected to the reaction flask, and the reaction mixture was deuterogenated with stirring for 16 h at room temperature. The resulting mixture was filtered to remove catalyst, and the filtrate was concentrated in vacuo. The residue was distilled under reduced pressure, thereby affording pure **7** (9.0 g, 85%): bp 95 °C (1 mm); IR (neat) 2245 (w), 2155 (w), 2135 (w), 2095 (w), 1735 (vs), 1445 (s), 1045 cm⁻¹ (s); ¹H NMR (CDCl₃) δ 3.66 (s); mass spectrum (70 eV), m/e (relative intensity) (no molecular ion), 119 (100.0), 118 (42.2), 117 (24.0), 91 (18.8), 59 (72.9), 58 (54.5), 57 (19.3), 32 (32.2), 21 (17.1).

1,4-Butanediammonium-1,1,4,4-d₄ Dinitrate (1c). A solution of **1a** (426 mg, 4.63 mmol) in 95 % aqueous ethanol (4 ml) was cooled at 0°C by application of an external ice bath. To this cold solution was added concentrated nitric acid (0.6 ml, 9.5 mmol) dropwise with vigorous stirring. The temperature of the reaction mixture was not permitted to exceed 0°C during the period of addition of nitric acid. After all of the nitric acid had been added, the ice bath was removed and the reaction mixture was permitted to warm to ambient temperature. After stirring for 10 minutes at room temperature, the reaction mixture was then filtered. The residue was washed with cold 95% ethanol and suction dried. The resulting material was recrystallized from ethanol-water mixed solvent, thereby affording pure **1c** (0.625 g, 62%) as a colorless microcrystalline solid: mp 140 °C, (lit⁷ mp 139 °C for non-deuterium labelled **1c**); IR (KBr pellet) 3048 (br, vs), 2400 (m), 1995 (m), 1759 (w), 1628 (m), 1520 (vw), 1354 (br, vs), 1073 (d, w), 1026 (m), 825 (m), 710 (m); ¹H NMR (DMSO-d₆) δ 1.6 (s, 4H), 5.2 (s, 6H); mass spectrum (70 eV), m/e (relative intensity) (no molecular ion), 74 (2.4), 73 (3.3), 62 (3.0), 61 (4.5), 57 (2.3), 46 (41), 44 (18), 32 (100), 30 (18).

1,4-Butanediammonium-2,2,3,3-d₄ Dinitrate (1d). The procedure described above for the preparation of **1c** was employed for the synthesis of **1d**. Pure **1d** was thereby obtained in 60% yield as an off-white microcrystalline solid: mp 139°C; IR (melt) 2350 (br, m), 1616 (w), 1517 (w), 1311 (br, vs), 1161 (vw), 1040 (vw), 826 (w), 722 (vw); ¹H NMR (DMSO-d₆) δ 2.3 (s, 4H), 7.9 (s, 6H); mass spectrum (70 eV), m/e (relative intensity) (no molecular ion), 75 (3.0), 74 (2.6), 73 (2.8), 63 (6.5), 62 (3.9), 47 (7.8), 46 (66.7), 45 (28.8), 44 (17.3), 43 (9.0), 31 (15.0), 30 (100.0).

SECTION IV

CONCLUSIONS

It has been demonstrated that the free bases 1a and 1b can be prepared in high purity by the non-aqueous methods described herein in relatively low, but reasonable yields. The free bases were reacted with concentrated nitric acid to provide their respective bis(ammonium nitrate) salts, 1c and 1d, in moderate yields. The NMR and mass spectra of the free bases and salts are consistent with the proposed structure and site-specific deuteration.

REFERENCES

1. Rogers, R. N., Janney, J. L. and Ebinger, M. H., Thermochim. Acta, Vol 59, page 237, 1982.
2. Shackelford, S. A., Beckmann, J. W., and Wilkes, J. S., J. Org. Chem., Vol 42, page 4201, 1977.
3. Shackelford, S. A., Coolidge, M. B., Goshgarian, B. B., Loving, B. A., Rogers, R. N., Janney, J. L. and Ebinger, M. H., J. Phys. Chem., Vol 89, page 3113, 1985, and references cited therein.
4. Mayerl, F. and Hesse, M., Helv. Chim. Acta, Vol 59, page 127, 1976.
5. Smith, R. G. and Daves, Jr., G. D., J. Org. Chem., Vol 43, page 2173, 1978.
6. Chambret, F. and Joly, D., Bull. soc. chim. France, page 1023, 1947.
7. McKenney, Jr., R. L., Struck, S. R., Hildreth, R. A. and Fryling, J. A., J. Energ. Mat. Vol 5, page 1, 1987.
8. Bennett, G. M. and Heathcoat, F., J. Chem. Soc., page 271, 1929.
9. Fried, S. and Richard, D. K., J. Am. Chem. Soc., Vol 62, page 3253, 1940.